

1096-34-133

Tracy Weyand* (weyand.tracy@gmail.com) and **Gregory Berkolaiko**. *Stability of Eigenvalues of Quantum Graphs*.

We consider the eigenvalues of the magnetic Schrödinger operator on a quantum graph as functions of the magnetic potential. We establish a simple relation between the Morse index of the magnetic eigenvalue and the number of zeros of the corresponding non-magnetic eigenfunction. This highlights an intricate relationship between zeros of an eigenfunction and the stability of the corresponding eigenvalue under magnetic perturbation.

In particular, let $\{\sigma_j\}_{j=1}^\beta$ be a set of generators of the fundamental group of a quantum graph Γ . The eigenvalues of the magnetic Schrödinger operator may be considered as functions of the magnetic flux $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_\beta)$ where $A(x)$ is the magnetic potential on Γ and

$$\alpha_i = \oint_{\sigma_j} A(x) dx.$$

Let ψ be the n -th eigenfunction of the ordinary Schrödinger operator (no magnetic potential) and assume that ψ is non-zero on the vertices of Γ . Let ϕ denote the number of internal zeros of ψ on Γ . We demonstrate that $(0, \dots, 0)$ is a non-degenerate critical point of $\lambda_n(\boldsymbol{\alpha})$ with Morse index equal to the nodal surplus of ψ , which is $\phi - (n - 1)$. (Received September 13, 2013)