

1116-05-179

**Ranjan Rohatgi\*** (rrohatgi@indiana.edu). *Lozenge tilings of halved hexagons with defects.*

MacMahon's boxed plane partition formula from over a century ago enumerates the lozenge tilings of a hexagon with side-lengths  $a, b, c, a, b, c$  in cyclic order on the triangular lattice. More recently several authors have enumerated the number of tilings for hexagons with different types of defects. Proctor treated the case where a "maximal staircase" is removed, and several others have found formulas for hexagons with triangles removed from the boundary. Potential lozenges used in the tiling of a region can be given weights. For a given tiling  $\mu$ , define  $\text{wt}(\mu)$  to be the product of the weights of all lozenges used in  $\mu$ . Rather than counting the number of tilings of a weighted region  $R$ , one tries to determine its *tiling generating function*,  $M(R)$  defined by  $M(R) = \sum_{\mu \in \mathcal{M}} \text{wt}(\mu)$  where  $\mathcal{M}$  is the set of all tilings of  $R$ . Evidently, if all weights are 1,  $M(R)$  is precisely the number of tilings of  $R$ . We present results for both unweighted and certain weighted hexagonal regions with both a maximal staircase and boundary triangles removed. Treating these regions as halved hexagons allows us, with the help of Ciucu's factorization theorem, to recover known results in a new way. (Received August 11, 2015)