

**Meeting:** 1003, Atlanta, Georgia, SS 35A, AMS-MAA Special Session on Tropical Geometry, I

1003-15-735      **Peter Butkovic\*** (p.butkovic@bham.ac.uk), School of Mathematics, The University of Birmingham, Edgbaston, B15 2TT Birmingham, England. *On the strong regularity of matrices in max-algebra.*

Let  $a \oplus b = \max(a, b)$  and  $a \otimes b = a + b$  for  $a, b \in \mathbb{R}$ . Extend  $(\oplus, \otimes)$  to matrices in the same way as in linear algebra.

Let  $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ . Then  $A$  is called *strongly regular* if there is a  $b \in \mathbb{R}^n$  such that the system  $A \otimes x = b$  has exactly one solution. If  $P_n$  is the set of all permutations of the set  $N = \{1, \dots, n\}$  and  $\pi \in P_n$  then the weight of  $\pi$  w.r.t.  $A$  is  $w(A, \pi) = \sum_{i \in N} a_{i, \pi(i)}$ .  $A$  is strongly regular if and only if there is exactly one permutation of maximal weight w.r.t.  $A$ . The set of all vectors  $b$  for which the system  $A \otimes x = b$  has one exactly solution is called the *simple image set* ( $S(A)$ ). The task of finding the simple image set can easily be transformed to the same question for normal matrices. If  $A$  is normal then  $S(A)$  is the interior of the eigenspace of  $A$ .

Let  $(G, \otimes, \leq)$  be a linearly ordered commutative group and define  $a \oplus b = \max(a, b)$ . The above criterion of strong regularity does not hold in general unless  $(G, \leq)$  is dense but an  $O(n^3)$  algorithm for checking the strong regularity in general groups exists. (Received October 02, 2004)